arXiv:1412.0105v1 [gr-qc] 29 Nov 2014

A note on Pathria's model of the universe as a black hole

S. Khakshournia *

Nuclear Science and Technology Research Institute (NSTRI), Tehran, Iran

December 2, 2014

Abstract

Pathria has shown that for the certain values of the cosmological constant, a pressureless closed Friedmann-Robertson-Walker universe can be the interior of a Schwarzschild black hole. We examine the Pathria's model from the point of view of the matching of the two different spacetimes through a null hypersurface. We first regard the event horizon of the black hole, which in the Pathria's model is identified with the radius of the universe at the point of its maximum expansion, as a null hypersurface separating the FRW interior from the vacuum Schwarzschild exterior, and then use the null shell formalism for it. It turns out that the matching is not smooth, and in fact, the null hypersurface is the history of a null shell admitting a surface pressure.

PACS numbers: 98.80.-k; 04.20.Jb

1 Introduction

Pathria [1] published a paper showing that a pressureless closed Friedmann-Robertson-Walker (FRW) universe can lie inside a Schwarzschild black hole existing in a bigger universe (see also [2]). He has first considered a FRW universe with positive 3D curvature, consisting of the dust matter, in a phase of decelerating expansion, with the value of the cosmological constant Λ such that $0 \leq \Lambda \leq \Lambda_c$, where $\Lambda_c = (2\pi\rho R^3)^{-2}$, with $\rho(t)$ and $R(t)$ are the energy density and the radius of curvature of the universe, respectively. In this case, the solution for the function $R(t)$ shows that in a later epoch it reaches its maximum value R_{max} increasing monotonically from the value $R_{max} = C$ ($C = \frac{8\pi}{3}\rho R^3$) to $R_{max} = \frac{3}{2}C$, as Λ increases from 0 to Λ_c. Pathria then compared this behavior with that of a 4D Schwarzschild black hole modified by the presence of the cosmological constant Λ and noted that the solutions for the radius of the event horizon R_s of the black hole are identically the same as for R_{max}, irrespective of the values of the parameters C and Λ, except that for $\Lambda > \Lambda_c$ neither R_s nor R_{max} exists [1]. He therefore suggested that the identity $R_{max} \equiv R_s$ is fundamental to the structure of the universe and concluded that at

*Email address: skhakshour@aeoi.org.ir

any epoch $R(t) \leq R_s$, meaning that the universe is indeed inside a black hole.
We note that this identification implies that the separating surface of the FRW interior from the vacuum exterior, while the universe reaches its maximum expansion, turns out to be (instantaneously) a null hypersurface. So what if we study the Pathria's cosmological model from the matching conditions point of view for the two different spacetimes through a null hypersurface? Hence, in this note, we formulate the general problem of gluing a FRW metric to a vacuum schwarzschild one along a null hypersurface. For this purpose, we use Barrabès-Israel null shell formalism [3] to investigate the matching conditions.
Conventions. Natural geometrized units, in which $G = c = 1$, are used throughout the paper. The null hypersurface is denoted by Σ. The symbol $|_\Sigma$ means "evaluated on the null hypersurface ". We use square brackets [F] to denote the jump of any quantity F across Σ. Latin indices range over the intrinsic coordinates of Σ denoted by ξ^a, and Greek indices over the coordinates of the 4-manifolds.

2 Matching Conditions

We consider a pressureless closed FRW expanding universe which can be described, in retarded time coordinates, by the following metric

$$ds_-^2 = -a^2 dv(2d\chi + dv) + a^2 \sin^2\chi(d\theta^2 + \sin^2\theta d\varphi^2), \tag{1}$$

where $a(v, \chi)$ is the scale factor, v is a null coordinate , and χ a spacelike coordinate which is constant along the comoving worldlines. The retarded time v is related to the cosmological time t_- by

$$dv = \frac{dt_-}{a} - d\chi. \tag{2}$$

The governing Friedmann equations are written as

$$\frac{\dot{a}^2}{a^2} + \frac{1}{a^2} - \frac{\Lambda}{3} = \frac{8\pi\rho}{3}, \tag{3}$$

$$\frac{\ddot{a}}{a} - \frac{\Lambda}{3} = -\frac{4\pi\rho}{3}, \tag{4}$$

where the dots denote the derivative with respect to t_-; Λ and ρ are the cosmological constant and the energy density, respectively.
Considering an external view of the universe, we assume the exterior of the FRW universe may be described by a Schwarzschild metric modified by the presence of the cosmological constant as follows

$$ds_+^2 = -du(f du + 2dr) + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \tag{5}$$

where

$$f = 1 - \frac{2M}{r} - \frac{\Lambda r^2}{3}, \tag{6}$$

with the Schwarzschild mass $M = 4\pi \int_0^r \rho r^2 dr = \frac{4\pi}{3}\rho r^3|_\Sigma$, being a constant for a dust-filled universe.

To match the FRW interior with the Schwarzschild exterior along the null hypersurface Σ situated at the maximum radius of the FRW expanding universe, we note that the requirement of the continuity of the induced metric on Σ yields the following matching conditions

$$\begin{aligned} r &\overset{\Sigma}{=} a\sin\chi, \\ 2d\chi &\overset{\Sigma}{=} -dv, \\ f du &\overset{\Sigma}{=} -2dr, \end{aligned} \tag{7}$$

where $\overset{\Sigma}{=}$ means that we should compute both sides of the equality at Σ. Taking $\xi^a = (u, \theta, \varphi)$ with $a = 1, 2, 3$ as the intrinsic coordinates on Σ, and u a parameter on the null generators of the hypersurface, we now calculate the tangent basis vectors $e_a = \partial/\partial\xi^a$ on both sides of Σ as

$$e^\mu_u|_- = \left(\frac{du}{d\chi}\right)^{-1}(-2,1,0,0)\,|_\Sigma, \qquad e^\mu_\theta|_- = \delta^\mu_\theta, \qquad e^\mu_\varphi|_- = \delta^\mu_\varphi, \tag{8}$$

$$e^\mu_u|_+ = \left(1, -\frac{f}{2}, 0, 0\right)|_\Sigma, \qquad e^\mu_\theta|_+ = \delta^\mu_\theta, \qquad e^\mu_\varphi|_+ = \delta^\mu_\varphi. \tag{9}$$

For further applications, using Eq. (7) together with (2) and (3), we can compute

$$\begin{aligned} \frac{du}{d\chi} &= \frac{du}{dr}\frac{dr}{d\chi}, \\ &= \frac{-2}{f}\left(-\frac{\partial a}{\partial\chi}\sin\chi + a\cos\chi\right)|_\Sigma, \\ &= \frac{-2}{f}\left(-a\dot{a}\sin\chi + a\cos\chi\right)|_\Sigma. \end{aligned} \tag{10}$$

Recalling that u is a parameter on the null geodesic generators of Σ, we choose the tangent-normal vector n^μ coinciding with the tangent basis vector associated with the parameter u, so that $n^\mu = e^\mu_u$ [4]. We may then complete the basis by a transverse null vector N^μ uniquely defined by the four conditions $n_\mu N^\mu = -1$, $N_\mu e^\mu_A = 0$ $(A = \theta, \varphi)$, and $N_\mu N^\mu = 0$. We find

$$N_\mu|_- = \frac{1}{2}\left(\frac{du}{d\chi}\right)(1,0,0,0)\,|_\Sigma, \tag{11}$$

$$N_\mu|_+ = (-1,0,0,0)|_\Sigma. \tag{12}$$

Furthermore, the induced metric on Σ, given by $g_{ab} = g_{\mu\nu}e^\mu_a e^\nu_b|_\pm$, is computed to be
$g_{ab} = \mathrm{diag}(0, r^2, r^2\sin^2\theta)\,|_\Sigma$, which is the same on both sides of the hypersurface. Defining a pseudo-inverse of the induced metric g_{ab} on Σ as $g^{ac}_* g_{bc} = \delta^a_b + n^a N_\mu e^\mu_b$, with $n^a = \delta^a_u$ [5], one gets $g^{ab}_* = \mathrm{diag}\left(0, \frac{1}{r^2}, \frac{1}{r^2\sin^2\theta}\right)|_\Sigma$.
The final junction conditions are formulated in terms of the jump in the transverse extrinsic curvature tensor across Σ. Using the definition $\mathcal{K}_{ab} = e^\mu_a e^\nu_b \nabla_\mu N_\nu$, we may therefore compute the transverse extrinsic curvature [5] on both sides of Σ. Its nonvanishing components on the minus side are found as

$$\begin{aligned} \mathcal{K}_{\theta\theta}|_- &= \sin^{-2}\theta\,\mathcal{K}_{\varphi\varphi}|_- \\ &= a\sin\chi\left(\frac{\cos^2\chi - \dot{a}^2\sin^2\chi}{f}\right)|_\Sigma, \\ &= a\sin\chi|_\Sigma, \end{aligned} \tag{13}$$

$$\begin{aligned}\mathcal{K}_{uu}|_{-} &= \frac{-1}{2}f_{,r} - 2\pi\rho a^3 f\left(\frac{dr}{d\chi}\right)^{-2}\sin\chi|_{\Sigma}, \\ &= \frac{-1}{2}f_{,r} + 2\pi\rho a|_{\Sigma},\end{aligned} \tag{14}$$

where we have used Eqs. (3), (4), (6), (7) and (10), and the term $\frac{dr}{d\chi}$ in Eq. (14) has been evaluated at $\chi = \frac{\pi}{2}$, corresponding to the maximum expansion of the FRW universe. The nonvanishing components of the transverse curvature on the plus side are

$$\mathcal{K}_{\theta\theta}|_{+} = \sin^{-2}\theta\mathcal{K}_{\varphi\varphi}|_{+} = r|_{\Sigma}, \tag{15}$$

$$\mathcal{K}_{uu}|_{+} = \frac{-1}{2}f_{,r}|_{\Sigma}. \tag{16}$$

Now Eqs. (13) and (15) show that the angular components of the transverse curvature are continuous across Σ. But from Eqs. (14) and (16) we infer the discontinuity of the "uu" component of extrinsic curvature across Σ, signaling the presence of a null shell of matter on the hypersurface Σ, with vanishing energy density, but a surface pressure which is computed as [5]

$$\begin{aligned}p &= -\frac{1}{8\pi}n^a n^b[\mathcal{K}_{ab}], \\ &= -\frac{1}{8\pi}[\mathcal{K}_{uu}], \\ &= \frac{\rho a}{4}|_{\Sigma}.\end{aligned} \tag{17}$$

The surface energy-momentum tensor of a lightlike shell is then written as $S^{ab} = pg_*^{ab}$, where the surface pressure p is measured by the observers comoving with the cosmological dust fluid and intersecting the hypersurface.
Therefore, the null hypersurface Σ, separating the closed FRW interior at the point of its maximum expansion from the vacuum Schwarzschild exterior, is the history of a null shell characterized by its surface pressure (17).

3 Conclusion

Applying the Barrabès-Israel null shell formalism, we have studied Pathria's cosmological model in which a pressureless closed FRW universe is entrapped within a Schwarzschild black hole, from the point of view of the matching conditions for the two different spacetimes glued together along a null hypersurface. As a result of the matching conditions, we see that the transition from the expansion to contraction in the FRW universe located on the event horizon of the Schwarzschild black hole can only be done through a null shell which is simply characterized by a surface pressure.

Note added in the published version: Very recently, Knutsen [6] critically examined Pathria's model from the point of view of smooth matching of a FRW interior region to the vacuum

Schwarzschild exterior spacetime along a timelike hypersurface. He points out that with the ill-employed notations in the Pathria's work, the matching conditions yield the unexpected results suggesting that Pathria's idea of the Universe as a black hole is untenable. However, it is worth addressing that in the limit when the timelike hypersurface separating the two spacetimes as studied in [6], is moved to the Schwarzschild horizon, it becomes a null hypersurface through which we have investigated the matching of the two spacetimes in the present work.

Acknowledgment

The author would like to thank the referee for helpful comments.

References

[1] R. K. Pathria, Nature **240**, 298 (1972).

[2] W. M. Stuckey, Am. J. Phys. **62**, 788 (1994); M. S. Berman, Astrophys. Space Sci. **311**, 359 (2007); N. J. Poplawski ; gr-qc/0901.0215.

[3] C. Barrabes and P. A. Hogan, *Singular Null Hypersurfaces in General Relativity*, (World Scientific, 2004).

[4] E. Poisson; gr-qc/0207101.

[5] C. Barrabès and W. Israel, Phys, Rev. D **43**, 1129 (1991).

[6] H. knutsen, Grav. Cosmol. **15**, 3, 273 (2009).